ELSEVIER

Journal of Immunological Methods 217 (1998) 201–202

JOURNAL OF IMMUNOLOGICAL METHODS

Contents of volume 217 (1998)

Numbers 1 and 2 – 1 August 1998

Research reports

$F(ab')_2$ molecules made from *Escherichia coli* produced Fab′ with hinge sequences conferring increased serum survival in an animal model
D.P. Humphreys, O.M. Vetterlein, A.P. Chapman, D.J. King, P. Antoniw, A.J. Suitters, D.G. Reeks, T.A.H. Parton, L.M. King, B.J. Smith, V. Lang and P.E. Stephens (UK) 1

A cytofluorometric method for the simultaneous detection of both intracellular and surface antigens of apoptotic peripheral lymphocytes
H. Lecoeur, E. Ledru and M.-L. Gougeon (France) 11

Bone marrow cellular composition in *Listeria monocytogenes* infected mice detected using ER-MP12 and ER-MP20 antibodies: a flow cytometric alternative to differential counting
M.F.T.R. De Bruijn, W. Van Vianen, R.E. Ploemacher, I.A.J.M. Bakker-Woudenberg, P.A. Campbell, W. Van Ewijk and P.J.M. Leenen (Netherlands, USA) 27

Detection of and discrimination between total and free human interleukin-4 and free soluble interleukin-4 receptor by ELISA
T. Jung, J.P.A. Bews, K.-H. Enssle, K. Wagner, C. Neumann and C.H. Heusser (Germany, Switzerland) 41

The matrix effects on kinetic rate constants of antibody–antigen interactions reflect solvent viscosity
C.L. Morgan, D.J. Newman, J.M. Burrin and C.P. Price (UK) 51

Apoptosis detection by annexin V binding: a novel method for the quantitation of cell-mediated cytotoxicity
Y. Shounan, X. Feng and P.J. O'Connell (Australia) 61

A modification of the JAM test is necessary for a correct determination of apoptosis induced by $FasL^+$ adherent tumor cells
C. Böhm, M.-L. Hanski, A. Gratchev, B. Mann, M.P. Moyer, E.-O. Riecken and C. Hanski (Germany, USA) 71

An ELISA to determine the biodistribution of human monoclonal antibody in tumor-xenografted SCID mice
K. Arai, K. Yoshinari, K. Matsumoto and H. Misaki (Japan) 79

A highly sensitive and specific assay using a novel human growth hormone cDNA reporter gene regulated by the human interleukin-4 inducible germline ε transcript promoter
C.-H. Jenh, M.A. Cox, D. Lundell, S.K. Narula and P.J. Zavodny (USA) 87

Establishment of the cells useful for murine interleukin-18 bioassay by introducing murine interleukin-18 receptor cDNA into human myelomonocytic KG-1 cells
M. Taniguchi, K. Nagaoka, S. Ushio, Y. Nukada, T. Okura, T. Mori, H. Yamauchi, T. Ohta, H. Ikegami and M. Kurimoto (Japan) 97

On-line flow displacement immunoassay for fatty acid-binding protein
W.A. Kaptein, J. Korf, S. Cheng, M. Yang, J.F.C. Glatz and R. Renneberg (China, Netherlands) 103

Inhibition of nonspecific binding of fluorescent-labelled antibodies to human eosinophils
S. Mahmudi-Azer, P. Lacy, B. Bablitz and R. Moqbel (Canada) 113

Removal of erythroid cells from umbilical cord blood mononuclear cell preparations using magnetic beads and a monoclonal antibody against glycophorin A
S.R. Elliott, P.J. Macardle and H. Zola (Australia) 121

Two high-affinity monoclonal IgG2a antibodies with differing thermodynamic stability demonstrate distinct antigen-induced changes in protein A-binding affinity
Z.I. Kravchuk, A.A. Chumanevich, A.P. Vlasov and S.P. Martsev (Belarus) 131

A simple and robust method for the complete dissociation of HIV-1 p24 and other antigens from immune complexes in serum and plasma samples
F. Steindl, C. Armbruster, K. Pierer, M. Purtscher and H.W.D. Katinger (Austria) 143

A comparative study of different methods for the assessment of apoptosis and necrosis in human eosinophils
G.M. Walsh, G. Dewson, A.J. Wardlaw, F. Levi-Schaffer and R. Moqbel (UK, Israel, Canada) 153

Optimization and comparison of the MTT assay and the ^{3}H-TdR assay for the detection of IL-2 in helper T cell precursor assays
C.A. Russell and L.L. Vindelov (Denmark) 165
A highly sensitive whole blood natural killer cell assay
K.V. Bromelow, J. Galea-Lauri, M.E. O'Brien and B.E. Souberbielle (UK) 177
Comparison of detection techniques for cytokine reverse transcriptase polymerase chain reaction; digoxigenin-labeled polymerase chain reaction permits sensitive detection of cytokine mRNA in rat heart allografts
J.G.M.C. Damoiseaux, R. Theunissen, C.P.M. Broeren, P.J.C. Van Breda Vriesman and A.M. Duijvestijn (Netherlands) 185

Short communication

A vector for the expression of recombinant monoclonal Fab fragments in bacteria
R. Burioni, P. Plaisant, F. Bugli, V. Delli Carri, M. Clementi and G. Fadda (Italy) 195

Contents of volume 217 (1998) 201

Notes to Authors 203

Protocol: Notes to Authors 205